

Quellen:

1. Du Y, Knopf H, Zhuang W et al. (2011) Pain perceived in a national community sample of German children and adolescents. *Eur J Pain* 15 (6): 649–657
2. Buksch J (2011) Health Behaviour in School-aged Children 2009/10 – a WHO Cross National Survey (HBSC). Sonderauswertung Universität Bielefeld
3. Perquin CW, Hazebroek-Kampschreur AA, Hunfeld JA et al. (2000) Pain in children and adolescents: a common experience. *Pain* 87 (1): 51–58
4. Hakala P, Rimpela A, Salminen JJ et al. (2002) Back, neck, and shoulder pain in Finnish adolescents: national cross sectional surveys. *BMJ* 325 (7367): 743
5. King S, Chambers CT, Huguet A et al. (2011) The epidemiology of chronic pain in children and adolescents revisited: a systematic review. *Pain* 152 (12): 2729–2738
6. Wiesel SW, Tsourmas N, Feffler HL, et al. A study of computer assisted tomography. The incidence of positive cat scans in an asymptomatic group of patients. *Spine* 1984;9:549-551
7. Boden SD, McCowin PR, Davis DO, et al. Abnormal magnetic-resonance scans of the cervical spine in asymptomatic subjects. A prospective investigation. *J Bone Joint Surg Am* 1990;72(8):1178-1184
8. Deyo R, Mirza S, Turner Y. Overtreating chronic back pain: time to back off? *J Am Board Fam Med* 2009;22(1):62-68
9. Scavone JC, Latshaw RF, Rohrar GV. Use of lumbar spine films: statistical evaluation at a teaching hospital. *JAMA* 1981;246:1105-1108
10. Simmons ED, Guyer RG, Graham-Smith A, et al. Radiograph assessment for patients with low back pain. *Spine J* 2003;3:S3-S5
11. Chou R, Fu R, Carrino J, et al. Imaging strategies for low back pain: systematic review and meta-analysis: *Lancet* 2009;373:463-472